

**Botanical Features and Anti-cancerous Potential of *Euphorbia nivulia* Buch.-Ham**

Muhammad Younus<sup>1,2\*</sup>, Muhammad M. Hasan<sup>2</sup>, Muhammad Hanif<sup>3</sup>, Ghulam Sarwar<sup>4</sup>, Khalil Ahmad<sup>5</sup>, Sajid M. ur Rehman<sup>1</sup>, Jafir H. Shirazi<sup>6</sup>, Qazi A. Jamil<sup>6</sup>, Kashif R. Khan<sup>7</sup>, Shahzada K. Syed<sup>8</sup>, Abuzar Ghaffari<sup>9</sup>, Muhammad S. Khan<sup>10</sup>, Muhammad Abdullah<sup>11</sup>, Muhammed A. Wazir<sup>12</sup>, Ayesha Rehman<sup>13</sup>

<sup>1</sup>Department of Pharmacognosy, Faculty of Pharmacy, Islamia University of Bahawalpur, Bahawalpur, Pakistan

<sup>2</sup>Department of Pharmacognosy, Faculty of Pharmacy and Pharmaceutical Sciences, University of Karachi, Karachi, Pakistan

<sup>3</sup>Department of Pharmacy, Benazir Bhutto Shaheed University, Lyari, Karachi, Pakistan

<sup>4</sup>Department of Botany, Faculty of Science, Islamia University of Bahawalpur, Pakistan

<sup>5</sup>University college of conventional medicines, Faculty of Pharmacy Islamia University of Bahawalpur, Pakistan

<sup>6</sup>Department of Pharmaceutics, Faculty of Pharmacy, The Islamia University of Bahawalpur, Pakistan

<sup>8</sup>Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Islamia University of Bahawalpur, Pakistan

<sup>9</sup>Islam College of Pharmacy, Sialkot, Pakistan

<sup>10</sup>Bahawalpur College of Pharmacy, BMDC, Bahawalpur, Pakistan

<sup>11</sup>Cholistan Institute of Desert Studies, The Islamia University of Bahawalpur, Bahawalpur, Pakistan

<sup>12</sup>Faculty of Pharmacy, University of Karachi, Karachi Pakistan

<sup>13</sup>Faculty of Pharmacy Islamia University of Bahawalpur, Bahawalpur, Pakistan

## ARTICLE INFO

## ABSTRACT

## Article history:

Received 07 May 2021

Revised 09 June 2021

Accepted 22 September 2021

Published online 02 October 2021

**Copyright:** © 2021 Younus *et al.* This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Euphorbia nivulia* – Buch.-Ham. (*En*), one of the members of Euphorbiaceae family, is rich in various phytoconstituents including flavonoids and polyphenolics. Many species of the genus *Euphorbia* have been reported to possess varying biological activities, including anti-proliferatory, cytotoxic, anti-inflammatory and antimicrobial activities. This medicinal plant species indigenous to the Cholistan desert of Bahawalpur, Pakistan, was evaluated for its botanical features and anti-cancerous potential. The species was evaluated for botanical features through anatomical evaluation and powder microscopy. Standard conventional procedures were employed for preliminary qualitative phytochemical screening of *En* (crude extract) and various fractions, named as *En* hexane (*En* hex), *En* chloroform (*En* ch), *En* butanol (*En* bt) and *En* aqueous (*En* aq) to identify various phytoconstituents. Toxicity to *Artemia salina* (Leach shrimp) larvae was assayed through Brine Shrimp Lethality Assay (BSLA) using Etoposide as standard. Anticancer activity of the plant against 3T3 and HeLa cell lines was assayed using the tetrazolium salt reduction assay. Doxorubicin and Cyclohexamide were used as standard. *En* showed optimum activity, 73.33% mortality against *Artemia salina*. Maximum cytotoxic activity against 3T3 and HeLa cells was shown by *En* and *En* hex. The current study may be helpful in proper identification of the plant species and suggests that *En* may be a suitable anticancerous candidate.

**Keywords:** *Euphorbia nivulia*, Botanical features, Anti-cancerous potentials, 3T3, HeLa cells.

## Introduction

Traditional medicines have been used for thousands of years in countries such as Egypt, India, China, and many other countries. Herbal products are still used as primary health care products in most third world countries.<sup>1</sup> Plants are an enormous reservoir of biologically active phytochemicals/compounds with diverse chemical structures and possessing protective/preventive properties against various diseases. Therefore, recent researches are devoted to the phytochemical investigation of higher plants with ethno-pharmacological relevance.<sup>2</sup> Today, the use of novel bioactive plant extracts in drug production for cancer prevention is dominating over synthetic chemicals.<sup>3,4</sup>

\*Corresponding author. E mail: [mkhan@iub.edu.pk](mailto:mkhan@iub.edu.pk)  
Tel: 0092 3464054083

**Citation:** Younus MY, Hasan MH, Hanif M, Sarwar G, Ahmad K, Rehman MS, Shirazi JH, Jamil OA, Khan KR, Syed SK, Ghaffari A, Khan MS, Abdullah M, Wazir MA, Rehman A. Botanical Features and Anti-cancerous Potential of *Euphorbia nivulia* Buch.-Ham. Trop J Nat Prod Res, 2021; 5(9):1591-1596. [doi.org/10.26538/tjnpr/v5i9.11](https://doi.org/10.26538/tjnpr/v5i9.11)

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria

Plant-derived biologically active compounds could act as protective agents against human carcinogenesis. Indo-Pakistan medicinal plants have a great history for their utility as a remedy for the treatment of a variety of ailments. The impact of medicinal plants and their isolated pure compounds is in a variety of therapeutic areas, but the major contribution is in the field of anti-infective and anticancer drug discovery.<sup>5</sup> In the past, plants have provided leads for the development of potential novel anti-cancerous agents like Taxol, paclitaxel, vinblastine and vincristine, etc., for a wide variety of leukemia, ovarian, breast and non-small-cell lung carcinomas. Screening of plant extracts has been of great interest to scientists to discover new anti-cancerous drugs.<sup>6</sup> Cancer is one of the most dangerous diseases in humans and presently demanding new anticancer agents from natural products. Over the last half-century most of the plant secondary metabolites and their derivatives have been used towards combating cancer.<sup>7,8</sup> Smoke, smoking, dietary imbalances, hormones, chronic infections/inflammations, plastic, artificial colors, etc., are the major causes of cancer. One of the major reasons of using herbs/plants as a source of anti-cancerous drug is that they are relatively simple, readily available, effective, economical and least toxic. To properly study the natural bioactive compounds of plant origin, development of certain bioassay techniques are necessary. In vitro methods allow the screening of large number of compounds for cytotoxicity against many types of cancer cell lines. It is not always possible to test

against cancer in animal models due to the system complexity. Therefore, *in vitro* assays are more sensitive to most anti-tumor agents than *in vivo* assays and usually requires less test material, time and money.<sup>10</sup> Crude plant extracts can be first assayed for particular activities and the active fractions then analyzed phytochemically.<sup>2</sup> A variety of cytotoxicity bioassays are now available.<sup>11</sup> Brine shrimp lethality assay (BSLA) is a general bioassay that appears capable of detecting a broad spectrum of bioactivity. This method provides a front-line screen that can be backed up by more specific and expensive bioassays, once the active compounds have been isolated. BSLA is predictive of cytotoxicity and pesticidal activity.<sup>12</sup> BSLA study cytotoxicity to reveal new anticancer compounds. Taxol™, an antitumor drug approved by FDA for treatment of ovarian, breast and non-small-cell lung carcinomas was discovered using the BSLA assay.<sup>13</sup> Ethno-pharmacological and scientific investigation has revealed anticancerous activity displayed by various members of *Euphorbia*, to justify the use of plants in ethno-medicine for the treatment of cancer and other various diseases. The Euphorbiaceae family, also known as the spurge family (the 4th largest family of the angiosperms), is composed of about 340 genera and over 8000 species,<sup>14</sup> which are distributed in several types of vegetation and habitats, mainly in the tropics. Approximately 195 species of one of its genus *Euphorbia* have been recorded from India.<sup>15</sup> *Euphorbia nivulia* – Buch.-Ham. is one of the members of the genus *Euphorbia* that belongs to the Euphorbiaceae family. It is rich in various phytoconstituents including flavonoids and polyphenolics. Preliminary phytochemical investigation of the plant showed the presence of phenolics and flavonoids as well as other several phytoconstituents.<sup>16</sup> Many species of the genus *Euphorbia* have been reported to possess varying biological activities, including anti-proliferation, cytotoxic, anti-inflammatory and antimicrobial activity, etc. These biological activities are attributed to the presence of specific classes of macro- and polycyclic diterpenes.<sup>17,18</sup> Ethnopharmacologically, a few species of the genus *Euphorbia* have been used for the treatment of skin diseases, gonorrhoea, migraine, intestinal parasites and as wart cures.<sup>19</sup> Generally, *Euphorbia* spp. has been used to treat asthma, leukemia, warts, as a laxative and diuretic, and have antiviral, antibacterial and antifungal activities.<sup>20-22</sup> As anticancer and antitumor, some *Euphorbia* species have shown activity against some cancers and tumors, like malignant melanoma, squamous cell carcinoma and lung cancer.<sup>23</sup> It has been reported that some species of *Euphorbia* have potential cytotoxic effect against different cell lines.<sup>24</sup> Presence of carcinogenic compounds, skin irritants, tumor promoters, anti-cancer agents and agents for overcoming multidrug resistance<sup>25</sup> is an incentive for detailed studies of other *Euphorbia* spp. Interest in natural products from *Euphorbia* species exhibiting biological activities was initiated with the isolation and structure elucidation of the tumour-promoting phorbol-12,13-diester of *Croton tiglium* L. (Euphorbiaceae) in 1968<sup>26</sup> and the description of the anti-tumour activity of a diterpene isolated from *Euphorbia esula* L. in 1976.<sup>27</sup> Release of ingenol mebutate (Picatos), a diterpene isolated from *Euphorbia peplus* L. used for the topical treatment of actinic keratosis<sup>28</sup> stimulated interest in natural products from *Euphorbia* species. Lathyrane diterpenoids isolated from *E. nivulia*, showed significant cytotoxic activity against Colo 205, MT2, and CEM cell lines.<sup>29</sup> The present work is an attempt to study the botanical features and anti-cancer activity of *Euphorbia nivulia*.

## Materials and Methods

### Plant material

Aerial parts (stem, branches, leaves and flowers) of fresh, well grown *Euphorbia nivulia* plant were collected in the months of March and April, 2015 from the premises of Hasilpur Road and adjoining areas of Bahawalpur region, Pakistan. The plant was authenticated by Ghulam Sarwar, a taxonomist in the Department of Botany, The Islamia University of Bahawalpur. A voucher specimen (EN-AP-05-12-041) was deposited in the herbarium of Pharmacology Research Lab, Department of Pharmacy, The Islamia University of Bahawalpur, Bahawalpur, Pakistan. The collected plant parts were chopped into

pieces and spread on filter paper under shade for drying at room temperature for forty days. Then, these were powdered by an electric grinder, and sieved through mesh No. 60. The powder was kept in closed amber coloured glass bottles. The fresh material was used for organoleptic evaluation, while the dried form was used for the physicochemical analysis and other parameters.

### Chemicals

All the chemicals and solvents used were of analytical grade and purchased from Sigma Aldrich, B. D. H. and Merck. Organic solvents (methanol/ethanol/acetone/water, DMSO), distilled water, test samples, *Artemia salina* (brine shrimps eggs), sea salt (38 g/L of D/W, pH 7.0, 3T3 (mouse fibroblast cells), HeLa cells, tetrazolium salt (obtained from Dr. Panjwani Center for Molecular Medicine and Drug Research and H.E.J Research Institute of Chemistry, University of Karachi, Karachi), hatching tray with perforated partition, lamp to attract brine-shrimp larvae, micro pipette (5, 50, 500 µL), vials tray, 9 vials samples for storage of *Artemia salina* eggs, flask, measuring cylinder, weighing balance, petri dish rotary evaporator, refrigerator, filter paper what-man # 1, binocular stereomicroscope (Labomad, USA).

### Botanical Evaluation

Botanical evaluation of the fresh aerial parts was performed as per WHO guidelines.<sup>30,31</sup>

### Microscopic evaluation

#### Powder microscopy

Small quantity of fine powder of the sample was placed on glass slide and 1 to 2 drops of mounting media were added. Digital binocular stereomicroscope (Labomed, USA) was used to observe the tissues and their organisations initially at 10x and 40x magnifications corresponding to low and high-power lens. The observed microscopic structures were identified by comparing with the previously reported standard.<sup>32,33</sup>

### Anatomical evaluation

Free hand transverse sections were prepared from leaves, stems and flowers of *Euphorbia nivulia*. Each specimen was placed on the glass slide and covered with cover slip, while binocular research microscope was used to observe the slide.<sup>34</sup> Safranin solution was used for the staining of thick-walled tissues like xylem, collenchyma, sclerenchyma cells, etc., while malachite green was used for the staining of thin-walled tissues like phloem, parenchyma cells, etc.

### Extraction and fractionation

Dried powdered plant material (10 kg) was macerated in 30 L of 70% aqueous ethanol at room temperature for 15 days with occasional stirring. The macerated mixture was filtered three times with muslin cloth separately and then further filtration was done by Whatman filter paper number 01. The filtrate was then evaporated under reduced pressure (-760 mmHg) and controlled temperature on the rotary evaporator (Heidolph Laborota 4000-efficient Germany and Buchi Rotavapor R-20). A thick, semisolid, and dark brown gummy mass was obtained which was then placed in the oven (Mettler Beschichtung Loading Model 100-800). The dried material was weighed, labeled and then stored at 4°C in an airtight container. The percentage yield was calculated. The condensed extract was used for further experiments. Moreover, n-hexane, chloroform and n-butanol was used in increasing order of polarity for partitioning of the extract into various fractions as previously described.<sup>31</sup>

### Preliminary phytochemical screening

Preliminary qualitative phytochemical screening of the *En* crude extract as well as various fractions, named as, *En* Hexane (*En* hex), *En* Chloroform (*En* ch), *En* Butanol (*En* bt) and *En* aqueous (*En* aq) fraction to identify the phytoconstituents like alkaloids, glycosides, flavonoids, tannins, saponins and phenols, etc., was carried out by using standard conventional procedures,<sup>16,30,31</sup> (detail of various tests performed is summarised in Table 1).

**Brine shrimp lethality assay**

The procedure for BSLA as described by Solis *et al.*<sup>35</sup> with little modification was adopted. Each sample (20 mg) was dissolved in 2.0 mL of ethanol and used to prepare the concentrations of 10, 100, 1000 µg/mL by transferring 5, 50 and 500 µL respectively to each vial. Three vials were made for each concentration, then the solvent was allowed to evaporate overnight.<sup>35</sup> Data was analyzed with Finney computer program to determine LD<sub>50</sub> values with 95% confidence intervals.<sup>36</sup>

**Cytotoxicity assay (3T3)**

Cytotoxic activity of phytoconstituents was evaluated in 96-well flat-bottomed micro plates by using the standard MTT (3-[4, 5-dimethylthiazole-2-yl]-2,5-diphenyl-tetrazolium bromide) colorimetric assay. MTT assay is potentially useful for assaying antiproliferative activities of materials, because of its specificity for living cells.<sup>37</sup> For this purpose, 3T3 (mouse fibroblast) cells were cultured in Dulbecco's Modified Eagle's Medium. The extent of MTT reduction to formazan within cells was calculated by measuring the absorbance at 540 nm, using a micro plate reader (Spectra Max plus, Molecular Devices, CA, USA). The cytotoxicity was recorded as concentration causing 50% growth inhibition (IC<sub>50</sub>) for 3T3 cells. The percent inhibition was calculated using following formula:

$$\% \text{ inhibition} = 100 - \left( \frac{\text{mean of O.D of test compound} - \text{mean of O.D of negative control}}{\text{Mean of O.D of positive control} - \text{mean of O.D of negative control}} \right) * 100$$

Results (% inhibition) were processed by using Soft- Max Pro software (Molecular Device, USA). The following concentrations were used for the calculations of IC<sub>50</sub>: 30 µM, 15 µM, 7.5 µM, 3.75 µM, 1.875 µM, 0.937µM, 0.468 µM and 0.234 µM

**Anticancer activity (HeLa cells)**

Cytotoxic activity of phytoconstituents was evaluated in 96-well flat-bottomed micro plates.<sup>37</sup> For this purpose, HeLa cells were cultured in Minimum Essential Eagle's Medium. The extent of MTT reduction to formazan within cells was calculated by measuring the optical density (O.D) absorbance at 570 nm, using a micro plate reader (Spectra Max plus, Molecular Devices, CA, USA). The cytotoxicity was recorded as concentration causing 50% growth inhibition (IC<sub>50</sub>) for HeLa. The percent inhibition was calculated by using following formula:

$$\% \text{ inhibition} = 100 - \left( \frac{\text{mean of O.D of test compound} - \text{mean of O.D of negative control}}{\text{Mean of O.D of positive control} - \text{mean of O.D of negative control}} \right) * 100$$

The results (% inhibition) were processed by using Soft- Max Pro software (Molecular Device, USA).

**Statistical Analysis**

Data were expressed as mean ± Standard Deviation (SD). Differences between mean values were analysed by one-way analysis of variance (ANOVA) with the level of significance set at P-value < 0.05.

**Results and Discussion**

Currently, different sophisticated modern research techniques and tools are available for the identification of drugs of plant origin, but macroscopic and microscopic methods are still one of the simplest, reliable, precise and economical methods to confirm the correct identity of the plant source. According to the World Health Organization,<sup>30</sup> the macroscopic and microscopic description of a plant is the first criteria to establish the identity and the degree of purity of material and should be done before any tests are conducted. Botanical /macroscopic evaluation is a qualitative test based on the study of morphological characters of crude drugs and serves as diagnostic tool. The current research was conducted on aerial parts of an indigenous medicinal plant of Cholistan Desert of Bahawalpur, Pakistan. Botanical features of *Euphorbia nivulia* reveal that the plant is a herb, 3-9 m in height. Stem is hard with straight, often whorled branches along with straight paired, stipulary spines often blackish. Leaves are 8.5-21.5 cm long and 4-5 cm wide. The fresh leaves are simple, fleshy, succulent, dark green with an herbaceous odour and characteristic taste; crowded at the end of branches, obviate oblong or spatulate and glabrous. Both dried and fresh leaves are cuneate shapes, the base is acute, and apex is sub-acute. Venation is pinnate. Lamina is entire and flat. The only midrib is visible in normal conditions. Secondary and tertiary veins are not visible due to their fleshy nature [Figure. 1(A,B,C,D,E)].

For microscopic study, fine powder mounted in chloral hydrate, glycerin and iodine revealed that the aerial parts contain fragments of epidermal tissues composed of thick-walled polygonal cells. Stomata are anomocytic type but infrequent. There are numerous rosette and needle-shaped calcium oxalate crystals and starch granules. Annular vessels showing pits on their secondary walls, and spiral vessels were also present. The cell walls of the fairly abundant (xylem) fibers are thickened or lignified. Multicellular, uniseriate, glandular trichomes are found scattered and attached to fragments of the epidermis with blunt ends [Figure. 2 (A, B, C, D, E, F)].

TS of the leaf showed the single-layered thick rectangular epidermal cells; epidermis is covered with cuticle. Cells of ground tissues are thin and spherical to irregular in shape. Laticifers are distributed in mesophyll region around the vascular bundles. Lamina is differentiated into mesophyll, upper and lower epidermis. On surface view both the epidermal cells appear polygonal in shape. Mesophyll tissue consists of thin-walled parenchyma cells differentiated into elongated palisade and spongy cells. Midrib is composed of bundle sheath, collenchyma and vascular bundles. Vascular bundles (VB) consist of lignified xylem (pink colour) whereas none lignified phloem (green colour) along with parenchyma. Calcium oxalate crystals were seen frequently among loosely arranged spongy mesophyll.

TS of young stem revealed normal dicot structures. Epidermis is thick, and below epidermis, there is a layer of collenchyma (Hypodermis), which is thick walled but still living. Secondary growth is extensive, and secondary phloem is distributed in distinct patches; the pith is large. Vascular bundles are composed of (pink) xylem and (green) phloem. The cortical region is rich in laticifers which secrete white thick milky latex.

**Table 1:** Phytochemical evaluation (*En cr* and fractions)


Phytochemical Class	ENH	ENC	ENB	ENA
Alkaloids	-	-	+	-
Glycosides	-	-	-	-
Tannins	-	-	+	+
Flavonoids	-	+	+	+
Saponins	-	-	+	+
Phenolic contents	-	+	+	+

+ = Present; - = Absent

TS of the flower showed that the outermost layer is epidermis. Below the epidermis ground tissues are composed of parenchyma cells. Inside ground tissues laticifers are present. Around central pith, vascular bundles are present [Figure. 3 (A, B, C, D)].

The microscopic studies of the powder and transverse sections showed the presence of various histological structures like epidermal cells, stomata, starch granules, vessels of different types, fibers, vascular bundles, laticifers and unicellular blunted and glandular trichomes.<sup>16</sup> The use of different stains or reagents helps to differentiate between various cellular structures based on their chemical composition.<sup>38</sup>


The current study aimed to evaluate the *in vitro* cytotoxic activity of hydro-alcoholic (70%) *En* and various fractions, named as *En* hexane, *En* chloroform, *En* butanol and *En* aqueous. The evaluation was performed using Brine Shrimp Lethality Assay (BSLA), 3T3 and HeLa cell lines in comparison to anticancer drugs. Results of BSLA are shown in Figure 4. The extract (*En*) showed the optimum activity against Brine Shrimp while its two fractions, i.e., *En* hexane and *En* butanol also showed activity. Etoposide was used as standard. Results of cytotoxicity assay (3T3) are shown in Figure 6. *En* showed good activity against 3T3 cells, while its fraction, *En* hexane also showed good activity. Doxorubicin was used as standard. Results of anticancer activity (on HeLa cells) are shown in Table 5. *En* showed good activity against HeLa cells while its fractions, *En* hexane also showed activity. Cyclohexamide was used as standard.




**Figure 3:** Anatomy A: TS Leaf (Upper Epidermis), B: TS Stem (Pith), C: TS Stem (Vascular Bundle), D: TS Flower (Pedicel)

Qualitative phytochemical screening tests were performed to detect phytochemicals in the crude extract and its fractions. Phytochemical analysis showed the presence of various phytochemicals like phenols, flavonoids, terpenoids, glycosides, alkaloids, saponin, steroids and tannins, etc. Brine shrimp lethality assay is commonly used in the primary screening of the crude extracts as well as fractions/isolated compounds to assess the toxicity and cytotoxic properties of the test samples. BSLA is used as a marker for general toxicity screening as it is a simple, efficient and inexpensive method for detection and isolation of insecticidal, pesticidal, cytotoxic, anti-neoplastic, antimalarial, and antifeedant compounds from plant extract, and is a suitable method to correlate the cytotoxic and other biological properties.<sup>39-41</sup> Many antitumor and pesticide natural products have been isolated using this bioassay.<sup>42, 43</sup> The variation in BSLA results (Table 1) may be due to the difference in the amount and kind of cytotoxic substances (e.g. tannins, flavonoids, triterpenoids, or coumarins) present in the crude extracts and fractions. Results showed decreased cell viability and cell growth inhibition in a dose-dependent manner. Furthermore, significant lethality of the crude extract (73.33% mortality) and *En* hexane and *En* butanol to brine shrimp indicates potent cytotoxic constituents. Cytotoxicity potentials of *En* against (3T3 and HeLa) cell line are due to various phytochemicals. Among these bioactive constituents, phenols, flavonoids, saponins, phenolic compounds and tannins, etc. possess strong antioxidant potential.


These antioxidants may prevent and cure cancer and other diseases by protecting the cells from damage caused by ROS/free radicals, the highly reactive oxygen compounds. Furthermore, phytochemicals including vitamins (A, C, E and K), carotenoids, terpenoids, flavonoids, polyphenol, tannins, saponins, pigments, enzymes and minerals display anticancer activities. These chemicals block various hormones and metabolic pathways associated with the development of cancer.<sup>44</sup> Several studies evaluated the relationships between the anti-proliferative activity of plant products and their phenolic content. A correlation exists between the structural oxidation state, position, number and nature of the substituents of the polyphenolic compounds and their anti-proliferative effects.<sup>45</sup> Phenolics may act as anti-proliferative agents to liquidate the tumor cells by disrupting the cell cycle.<sup>46</sup> Moreover, triterpenes and saponins may contribute to strong anticancer activities.<sup>47</sup> Similarly, some of steroidal saponins had antimicrobial and cytotoxic activity.<sup>48,49</sup> Hydroxycoumarins are typical phenolic compounds that are made up of fused benzene and a-pyrone rings, and possess antioxidant, hepatoprotective, antiviral, and anticarcinogenic activities, therefore act as potent metal chelators and free radical scavengers.<sup>50</sup> Recent reports show that tannins may have potential value as cytotoxic and/or antineoplastic agents.<sup>51</sup>




**Figure 1:** *Euphorbia nivulia* A: Plant habit, B: Phyllotaxy, C: Fresh leaves, D: Stem with black spines, E: Bark & Wood (TS Stem)




**Figure 2:** Powder Study A: Parenchyma cells, B: Calcium oxalate crystals, C: Trichomes, D: Spiral vessel & fiber, E: Pitted vessel, F: Mesophyll



**Figure 4:** Lethality of various fractions (1000  $\mu\text{g/mL}$ ) of *E. nivulia* on Brine Shrimp eggs against Etoposide



**Figure 5:** Cytotoxicity studies of various fractions (30  $\mu\text{g/mL}$ ) of *E. nivulia* on Hela cells against Doxorubicin



**Figure 6:** Cytotoxicity studies of various fractions (30  $\mu\text{g/mL}$ ) of *E. nivulia* on 3T3 cells against Cyclohexamide

## Conclusion

The current study is a step towards the identification of the plant species, i.e., *Euphorbia nivulia* (En) belonging to Cholistan Desert of Pakistan. The present study also revealed that En is very rich in various phytochemicals including polyphenolic compounds like phenolic acids and flavonoids with excellent antioxidant properties. Moreover, the study recommends that En may be a good candidate as anticancerous agent for developing new formulation(s).

## Conflict of Interest

The authors declare no conflict of interest.

## Authors' Declaration

The authors hereby declare that the work presented in this article is original and any liability for claims relating to the content of this article will be borne by them.

## Acknowledgements

Authors are thankful to Department of Pharmacognosy, Faculty of Pharmacy, The Islamia University of Bahawalpur, Bahawalpur, Pakistan; and Department of Pharmacognosy, Faculty of Pharmacy and Pharmaceutical Sciences, University of Karachi, Pakistan, for providing lab facilities to accomplish this work.

## References

- Mans DR, Da Rocha AB, Schwartsmann G. Anti-cancer drug discovery and development in Brazil: targeted plant collection as a rational strategy to acquire candidate anti-cancer compounds. *The Oncol.* 2000; 5(3):185-198.
- Harborne AJ. *Phytochemical methods a guide to modern techniques of plant analysis.* springer science & business media; 1998. 55 p.
- Wu J, Wu Y, Yang BB. Anticancer activity of *Hemsleya amabilis* extract. *Life Sci.* 2002; 71(18):2161-2170.
- Kwiecinski MR, Felipe KB, Schoenfelder T, de Lemos Wiese LP, Rossi MH, Gonalez E, Felicio JD, Wilhelm Filho D, Pedrosa RC. Study of the antitumor potential of *Bidens pilosa* (Asteraceae) used in Brazilian folk medicin. *J Ethnopharmacol.* 2008; 117(1):69-75.
- Newman DJ and Cragg GM. Natural products as sources of new drugs over the last 25 years. *J Nat Prod.* 2007; 70(3): 461-477.
- Dimayuga RE and Garcia SK. Antimicrobial screening of medicinal plants from Baja California Sur, Mexico. *J Ethnopharmacol.* 1991; 31(2):181-192.
- Newman DJ, Cragg GM, Snader KM. The influence of natural products upon drug discovery. *Nat Prod Rep.* 2000; 17(3): 215-234.
- Butler MS. The role of natural product chemistry in drug discovery. *J Nat Prod.* 2004; 67(12):2141-2153.
- El-Menshawi BS, Fayad W, Mahmoud K, El-Hallouty SM, El-Manawaty M. Screening of natural products for therapeutic activity against solid tumors. *Indian J Exp.* 2010; 48(3):258-264.
- Cragg GM, Boyd MR, Khanna R, Kneller R, Mays TD, Mazan KD, Newman DJ, Sausville EA. International collaboration in drug discovery and development: the NCI experience. *Pure Appl Chem.* 1999; 71(9): 1619-1633.
- Hostettmann K. *Assays for bioactivity.* Academic Press, London. 1991. 25-26 p.

12. Ghisalberti EL. Detection and isolation of bioactive natural products. In: *Bioactive Natural Products*. CRC Press; 2007. 25-90 p.
13. He L, Orr GA, Horwitz SB. Novel molecules that interact with microtubules and have functional activity similar to Taxol™. *Drug Discov Today*. 2001; 6(22): 1153-1164.
14. Younus M, Hasan MM, Ali S, Saddiq B, Sarwar G, Ullah MI, Maqsood A, Ahmar S, Mora-Poblete F, Hassan F, Chen JT. Extracts of *Euphorbia nivulia* Buch.-Ham. showed both phytotoxic and insecticidal capacities against *Lemna minor* L. and *Oxycarenus hyalinipennis* Costa. *PLOS ONE*. 2021; 16(4):e0250118.
15. Aditya S. A revision of geophytic *Euphorbia* species from India. *Euphorbia World*. 2010; 6(1):18-21.
16. Younus M, Hasan MM, Abbas K, Sarwar G. Pharmacognostic and physicochemical screening of *Euphorbia nivulia* Buch.-Ham. *Pak J Pharm Sci*. 2019; 32(3):1111-1119.
17. Trease GE and Evans WC. *Pharmacognosy*. (13<sup>th</sup>). ELBS/Bailliere Tindall, London; 1989. 345-346 p.
18. Cateni F, Falsone G, Zilic J. Terpenoids and glycolipids from Euphorbiaceae. *Mini Rev Med Chem*. 2003; 3(5):425-437.
19. Singla AK and KamLa P. Phytoconstituents of *Euphorbia* species. *Fitoter*. 1990; 41(6): 483-516.
20. Vijaya K, Ananthan S, Nalini R. Antibacterial effect of theaflavin, polyphenon 60 (*Camellia sinensis*) and *Euphorbia hirta* on *Shigella* spp.—a cell culture study. *J Ethnopharmacol*. 1995; 49(2):115-118.
21. Betancur-Galvis LA, Morales GE, Forero JE, Roldan J. Cytotoxic and antiviral activities of Colombian medicinal plant extracts of the *Euphorbia* genus. *Memórias do Instituto Oswaldo Cruz*. 2002; 97(4):541-546.
22. Yu FR, Lian XZ, Guo HY, McGuire PM, Li RD, Wang R, Yu FH. Isolation and characterization of methyl esters and derivatives from *Euphorbia kansui* (Euphorbiaceae) and their inhibitory effects on the human SGC-7901 cells. *J Pharm Pharm Sci*. 2005; 8(3):528-535.
23. Sreenika G, Naga SK, Lakshmi BV, Thulja P, Sudhakar M. Antioxidant and antitumor activity of *Euphorbia milii* flower extract against in vivo breast cancer and colon cancer in mice. *World J Pharm Pharm Sci*. 2015; 4: 912-934.
24. Wang ZY, Liu HP, Zhang YC, Guo LQ, Li ZX, Shi XF. Anticancer potential of *Euphorbia helioscopia* L extracts against human cancer cells. *Anat Rec*. 2012; 295(2):223-233.
25. Sadeghi-Aliabadi H, Sajjadi SE, Khodamoradi M. Cytotoxicity of *Euphorbia macroclada* on MDA-MB-468 breast cancer cell line. *Iran J Pharm Sci*. 2009; 5(2): 103-108.
26. Corea G, Fattorusso E, Lanzotti V, Di Meglio P, Maffia P, Grassia G, Ialenti A. & Ianaro A. Discovery and biological evaluation of the novel naturally occurring diterpene pepluanone as antiinflammatory agent. *J Med Chem*. 2005; 48(22):7055-7062.
27. Hecker E. Cocarcinogenic principles from the seed oil of *Croton tiglium* and from other Euphorbiaceae. *Cancer Res*. 1968; 28(11):2338-2348.
28. Berman B. New developments in the treatment of actinic keratosis: focus on ingenol mebutate gel. *Clin Cosmet Investig Dermatol*. 2012; 5:111-122.
29. Ravikanth V, Reddy VL, Reddy AV, Ravinder K, Rao TP, Ram TS, Kumar KA, Vamanarao DP, Venkateswarlu Y. Three new ingol diterpenes from *Euphorbia nivulia*: evaluation of cytotoxic activity. *Chem Pharm Bull*. 2003; 51(4):431-434.
30. World Health Organization. Quality control methods for medicinal plant materials. World Health Organization; 1998. 187 p.
31. Brain KR and Turner TD. The practical evaluation of phytopharmaceuticals. Bristol: Wright-Scientechica; 1975. 81 p.
32. Jackson BP and Snowdon DW. Atlas of microscopy of medicinal plants, culinary herbs and spices. Belhaven Press; 1990. 75 p.
33. Youngken HW. Textbook of Pharmacognosy. (6th ed). The Blakiston Company, Philadelphia. 205-206 p.
34. Ruzin SE. Plant microtechnique and microscopy. New York: Oxford University Press; 1999. 90 p.
35. Solis PN, Wright CW, Anderson MM, Gupta MP, Phillipson JD. A microwell cytotoxicity assay using *Artemia salina* (brine shrimp). *Planta Med*. 1993; 59(3):250-252.
36. Wiegand H and Finney DJ. Probit analysis. Aufl. Cambridge University Press, Cambridge; 1971. 80 p.
37. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods*. 1983; 65(1-2):55-63.
38. Mali PY and Panchal SS. Pharmacognostical and physicochemical standardization of *Euphorbia nerifolia* leaves. *Pharmacogn Mag*. 2017; 9(5):696-705.
39. Montanher AB, Pizzolatti MG, Brighente IM. An application of the brine shrimp bioassay for general screening of Brazilian medicinal plants. *Acta Farm Bonaerense*. 2002; 21(3):175-178.
40. Lalian JA, Nuñez OM, Uy MM. Brine shrimp (*Artemia salina*) bioassay of the medicinal plant *Pseudelephantopus spicatus* from Iligan City, Philippines. *Int Res J Biol Sci*. 2014; 3: 47-50.
41. Olowa LF and Nuñez OM. Brine shrimp lethality assay of the ethanolic extracts of three selected species of medicinal plants from Iligan City, Philippines. *Int Res J Biol Sci*. 2013; 2(11): 74-77.
42. McLaughlin JL, Chang CJ, Smith D. Bench-top bioassays for the discovery of bioactive natural products: an update. *Stud Nat Prod Chem*. 1991; 9:383-409.
43. Meyer BN, Ferrigni NR, Putnam JE, Jacobsen LB, Nichols DE, McLaughlin JL. Brine shrimp: a convenient general bioassay for active plant constituents. *Planta Med*. 1982; 45(5):31-34.
44. Prajapati ND, Purohit SS, Sharma AK, Kumar T. A Hand Book of Medicinal Plants. (1st ed). Agrobios Publisher Jodhpur, India; 2003. 76 p.
45. Yanez J, Vicente V, Alcaraz M, Castillo J, Benavente-Garcia O, Canteras M, Teruel JA. Cytotoxicity and antiproliferative activities of several phenolic compounds against three melanocytes cell lines: relationship between structure and activity. *Nutr Cancer*. 2004; 49(2):191-199.
46. Conforti F, Ioele G, Statti GA, Marrelli M, Ragno G, Menichini F. Antiproliferative activity against human tumor cell lines and toxicity test on Mediterranean dietary plants. *Food Chem Toxicol*. 2008; 46(10): 3325-3332.
47. Dzhambazov B, Daskalova S, Monteveva A, Popov N. In vitro screening for antitumor activity of *Clinopodium vulgare* L.(Lamiaceae) extracts. *Biol Pharm Bull*. 2002; 25(4): 499-504.
48. Zhang C, Feng S, Zhang L, Ren Z. A new cytotoxic steroidal saponin from the rhizomes and roots of *Smilax scobinicaulis*. *Nat Prod Res*. 2013; 27(14):1255-1260.
49. Xu J, Feng S, Wang Q, Cao Y, Sun M, Zhang C. Four new furostanol saponins from the rhizomes and roots of *Smilax scobinicaulis* and their cytotoxicity. *Molecules*. 2014; 19(12):20975-20987.
50. Kostova I. Synthetic and natural coumarins as cytotoxic agents. *Anti-Cancer Agents Med*. 2005; 5(1):29-46.
51. Claustra AL, Madulid RS, Aguinaldo AM, Espeso EI, Guevara BQ, Nonato MG, Santos MA, de Castro-Bernas G, Gonzales RE, del Castillo-Solevilla RC, Ysrael MC. A Guidebook to Plant Screening: Phytochemical and Biological. University of Santo Tomas Publishing House, Espana, Manila; 2005. 102-05 p.