

***In vitro* Inhibitory HMG-CoA Reductase Activity of Purified Polyphenol Compounds from Water Lettuce (*Pistia Stratiotes*) Leaf Extract**

Sabri Sudirman*, Miftahul Janna, Herpandi, Indah Widiastuti

Fisheries Product Technology, Faculty of Agriculture, Universitas Sriwijaya, Indralaya 30862, Ogan Ilir Regency, South Sumatra, Indonesia

ARTICLE INFO

ABSTRACT

Article history:

Received 02 June 2022

Revised 28 June 2022

Accepted 23 July 2022

Published online 03 August 2022

Copyright: © 2022 Sudirman *et al.* This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

The 3-hydroxy-3-methylglutaryl-coenzyme-A (HMG-CoA) reductase is an enzyme that plays a role in the synthesis of cholesterol. The side effect of synthetic anti-cholesterol drugs has led to the demand for natural HMG-CoA reductase inhibitors such as plant extract. Therefore, this study aimed to determine the inhibitory HMG-CoA reductase activity of polyphenol compounds from water lettuce (*Pistia stratiotes*) leaf extract. The research was conducted using the crude and purified extracts of water lettuce. The total polyphenols, flavonoids, and HMG-CoA reductase inhibitory activity assay were carried out by *in vitro* analysis. The values obtained were analyzed quantitatively, followed by the use of an independent sample t-test and presented in graphical form. The results showed that the total polyphenols in each crude and purified extracts were 29.03 mg GAE/g dry sample and 65.63 mg GAE/g dry sample, respectively, while the total flavonoids were 27.58 mg QE/g dry sample and 88.02 mg QE/g dry sample, respectively. The inhibitory activity of the HMG-CoA reductase enzyme showed that the purified extract showed percentage inhibition of 34.74% which was higher than that of the crude extract which was 2.61%. This indicated that the purified extract of water lettuce has higher levels of polyphenols and flavonoids that can inhibit the HMG-CoA reductase enzyme more effectively than the crude extract.

Keywords: Anti-hypercholesterolemia, HMG-CoA reductase, Polyphenol, Water lettuce.

Introduction

Currently, society's lifestyle is experiencing significant changes such as high consumption of fast food and lack of physical activity including exercise. This condition usually leads to a metabolic imbalance in the body that can cause the accumulation of fat and increase cholesterol. A previous study reported that cholesterol is a very important compound in human life. However, high cholesterol is closely linked with many other medical problems, especially elevated levels of low-density lipoprotein-cholesterol (LDL-C).¹ It was also discovered that high cholesterol levels or hypercholesterolemia can cause cardiovascular diseases (CVDs), such as atherosclerosis, stroke, and heart disease.²

Pharmacological management such as the consumption of statin drugs is a treatment for reducing cholesterol levels in the body by inhibiting the 3-hydroxy-3-methylglutaryl coenzyme-A (HMG-CoA) reductase enzyme. The HMG-CoA reductase is an important enzyme that is involved in cholesterol synthesis.³ However, the use of this drug has shown some adverse effects, which include headache, muscle pain, and digestive system problems.^{4,5} Therefore, investigations are being carried out using plant extract as a functional or food supplement to discover an alternative inhibitor of the HMG-CoA reductase.

Water lettuce (*Pistia stratiotes*) is an aquatic plant that contains some bioactive compounds, such as polyphenols, flavonoids, and tannins.^{6,7} A previous study also reported polyphenol compounds show lowering cholesterol activity by inhibiting the HMG-CoA reductase enzyme.⁸

*Corresponding author. E mail: sabrisudirman@unsri.ac.id
Tel: +62 711580934

Citation: Sudirman S, Janna M, Herpandi, Indah Widiastuti I. *In vitro* Inhibitory HMG-CoA Reductase Activity of Purified Polyphenol Compounds from Water Lettuce (*Pistia Stratiotes*) Leaf Extract. Trop J Nat Prod Res. 2022; 6(7):1131-1134. doi.org/10.26538/tjnpr/v6i7.15

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria.

It was also reported that flavonoid possess the potential anti-cholesterol activity and prevent CVDs.⁹ Since polyphenol extraction with organic solvent was obtained as a crude extract composed of other non-polyphenol components, such as lipid, sugar, and organic acids, a purification process is required to remove these compounds.¹⁰ Based on this condition, the authors hypothesized that the polyphenol extract showed different activity before and after purification. Therefore, this study aimed to investigate the polyphenol content from water lettuce (*Pistia stratiotes*) and its HMG-CoA reductase activity.

Materials and Methods*Sample preparation and extraction*

The water lettuce (*Pistia stratiotes*) was collected in September 2021 from Sukaraja Village, South Sumatra, Indonesia. The sample collected was authenticated at Microbiology and Biotechnology Laboratory of Fisheries Product Technology, Universitas Sriwijaya (FPT0015092022). The leaf was rinsed with distilled water and oven-dried at 45°C until constant weight is reached. After the drying process, the sample was milled to a size of 40 mesh using a grinding machine (Microphyte disintegrator B-One DM-120M) and kept for the extraction process. The polyphenol compound was extracted by the maceration method using 70% ethanol as a solvent at room temperature for 3 hours.⁷ Briefly, 20 mg of sample and 200 mL of 70% ethanol were mixed in the Erlenmeyer flask, then stirred using a magnetic stirrer. After 3 hours, the filtrate and residue were separated with a filter paper (Whatman No. 42). The filtrate was kept in a new collection tube and the residue was extracted by a fresh solvent under the same condition as the first extraction, which was carried out five repetitions. The filtrate-mixed was evaporated using a vacuum rotary evaporator at 40°C to obtain the concentrated extract. Half of the concentrated extract was dried using a freeze dryer to obtain polyphenol extract in powder form (crude extract). Meanwhile, purified extract was obtained through a HyperSep Retain PEP cartridge.

Purification process

The purification process was performed by solid-phase extraction (SPE) and a HyperSep Retain PEP cartridge (Part. No. 60107-212, Thermofisher Scientific) as described by Pérez-Magariño *et al.*¹¹ Briefly, 2 mL of distilled water and then 2 mL of methanol was rinsed for cartridge preconditioned. Subsequently, 2 mL of crude extract was loaded into the cartridge and the sample was eluted by using 2 mL of n-hexane and 2 mL of 1 N H₂SO₄. The cartridge was washed with absolute methanol to obtain purified polyphenol extract in an aqueous form and dried with a freeze dryer to obtain the powder form of purified extract.

Total polyphenol and flavonoid analysis

The total polyphenol and flavonoid contents were analyzed according to Chandra *et al.*¹² Moreover, total polyphenol content was analyzed using Folin-Ciocalteu's phenol reagent. Briefly, 50 mg of dried extract was dissolved in 10 mL of methanol. Then 0.2 mL of this solution was mixed with 0.2 mL Folin-Ciocalteu's phenol reagent. After 5 min, the mixture was added with 1 mL of 8% sodium carbonate and then volume up to 3 mL with distilled water. The mixture was incubated at room temperature and in dark conditions for 30 min. Subsequently, the absorbance was measured at 765 nm by a spectrophotometer (Genesys 150 ThermoScientific, Massachusetts, USA). The gallic acid (GA) was used as a standard and the total polyphenol content (TPC) was expressed as mg gallic acid equivalent (GAE) per g of dry sample.

Total flavonoid content was analyzed using the aluminum chloride method. Briefly, 50 mg extract was dissolved in 10 mL of methanol. Then, 1 mL of extract solution was mixed with 1 mL of 2% aluminum chloride and allowed to react at room temperature for 60 min. After the reaction time, 1 mL of mixture was made up to 10 mL and kept for 5 min. After that, the absorbance was measured at 420 nm by a spectrophotometer (Genesys 150 ThermoScientific, Massachusetts, USA). The quercetin was used as a standard and total flavonoid

content (TFC) was expressed as mg quercetin equivalent (QE) per g of dry sample.

HMG-CoA reductase inhibitory activity assay

The 3-hydroxy-3-methylglutaryl Coenzyme A (HMG-CoA) reductase inhibitory activity was measured using commercial kits from Sigma-Aldrich Co. (CS1090-1KT, Merck) and performed according to the manufacturers' protocol. Briefly, 50 mg of each crude and purified extract was dissolved in assay buffer. The solution was centrifuged at 3,000 rpm for 5 min, filtrated and the supernatant was kept for the next steps. Then, 5 µL of each extract and 0.5 µL of pravastatin were pipetted into reaction tubes according to the Reaction Mixed (Table 1). The reaction was observed at 340 nm by using a spectrophotometer every 20 s for 10 min. The HMG-CoA reductase activity and percentage of inhibitor were calculated according to these formulas:

$$= \frac{(\Delta A_{340}/\text{min}_{\text{sample}} - \Delta A_{340}/\text{min}_{\text{blanko}}) \times \text{TV}}{12.44 \times V \times 0.6 \times \text{LP}} \quad \text{Enzyme activity (Units/mgP)}$$

Where: TV = Total volume of the reaction in ml (1 ml for cuvettes and 0.2 ml for plates); V = volume of enzyme used in the assay (ml); LP = Light path in cm (1 for cuvettes).

$$\text{Inhibition (\%)} = \frac{\text{Enzyme activity without sample} \left(\frac{\text{Unit}}{\text{mgP}}\right) - \text{Enzyme activity with sample} \left(\frac{\text{Unit}}{\text{mgP}}\right)}{\text{Enzyme activity without sample} \left(\frac{\text{Unit}}{\text{mgP}}\right)} \times 100\%$$

Statistical analysis

All data are expressed as the mean ± standard deviation (SD) and analyzed by one-way ANOVA with Duncan's post-hoc test ($p < 0.05$) using SPSS (v.22.0; IBM Corp., Armonk, NY, USA).

Table 1: Reaction Mixture of HMG-CoA reductase inhibitory activity (for 1 mL of cuvette)

Sample	Assay buffer	Crude extract (CE)	Purified extract (PE)	Pravastatin (Prava)	NADPH	HMG-CoA	HMG-CoA Reductase
Blanko	920 µL	-	-	-	20 µL	60 µL	-
HMG-CoA reductase activity	915 µL	-	-	-	20 µL	60 µL	5 µL
CE inhibition	910 µL	5 µL	-	-	20 µL	60 µL	5 µL
PE inhibition	910 µL	-	5 µL	-	20 µL	60 µL	5 µL
Prava inhibition	910 µL	-	-	5 µL	20 µL	60 µL	5 µL

Results and Discussion

Total polyphenol and flavonoid content

The results showed that the purified extract has significantly higher total polyphenol and flavonoid contents compared to the crude extract as shown in Figure 1. Total polyphenol contents (TPC) of the crude and purified extracts were about 29.03 ± 4.15 mg GAE/g dry sample and 65.63 ± 6.51 mg GAE/g dry sample. Meanwhile, total flavonoids were about 27.58 mg QE/g dry sample and 88.02 mg QE/g dry sample, respectively. It was also found that the purified extract is majorly composed of polyphenol and flavonoid compounds due to unwanted compounds are being removed during the purification process. A previous study reported that the quality or bioactivity of crude extract was increased after purification.¹³ This is because unwanted and undesirable substances have been removed, thereby improving the overall quality of the crude extract.¹⁴ A solid-phase extraction (SPE) method was used for the purification process due to its rapid, simple, and economic.¹⁵ The n-hexane was applied for the removal of non-polar from the crude extract and a low concentrate of sulfuric acid was used to remove non-polyphenol polar compounds, such as sugar and organic acids.¹¹ A previous study reported that the total polyphenol of crude extract of *Inga edulis* increased from 496.5 mg GAE/g to 518.8 mg GAE/ after purification process.¹⁶ Furthermore, the luteolin (a flavonoid compound) increased from 0.68

mg/g to 3.52 mg/g and some flavonoids compounds of olive pomace also improved after the purification process.¹⁷

HMG-CoA reductase inhibitory activity

Figure 2 showed the representative reduction of absorbance at 340 nm during the measurement of the HMG-CoA reductase activity assay. After the calculation, it was found that purified extract with a value $34.74 \pm 5.40\%$ showed HMG-CoA reductase activity significantly higher than crude extract ($2.61 \pm 0.28\%$) as shown in Figure 3.

The increased inhibition of purified extract to HMG-CoA reductase due to it composed of high concentrations of polyphenol and flavonoid compounds. A previous study reported that polyphenols and flavonoids from Malabar spinach (*Basella alba*) leaf have the ability to inhibit HMG-CoA reductase activity.¹⁸ A previous reference also reported that isoflavon (a flavonoid compound) inhibits HMG-CoA reductase activity as an inhibitor competitive during cholesterol synthesis.¹⁹ According to Chen *et al.*, catechin (a polyphenol compound) successfully reduced cholesterol levels through *in vitro* and *in vivo* experiments.²⁰ Additionally, Islam *et al.*⁸ reported that these compounds can block the electron transfer on the substrate HMG-CoA and bind on HMG-CoA reductase on the NADP⁺ binding site. The HMG-CoA reductase is an important enzyme that is involved in cholesterol synthesis. This enzyme catalyzed HMG-CoA to

Coenzyme A and mevalonate. The mevalonate then was converted to cholesterol by the mevalonate pathway.³ Therefore, inhibited HMG-CoA reductase is an effective way to reduce cholesterol levels in human and animal experiments.²¹

lettuce (*Pistia stratiotes*) has a potential to be developed as an alternative anti-hypercholesterolemia agent.

Conflict of Interest

The authors declare no conflict of interest.

Authors' Declaration

The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.

Acknowledgements

This work was financially supported by the Universitas Sriwijaya within the framework of the Indonesia Directorate General of Higher Education, Research and Technology Project, Ministry of Education, Culture, Research, and Technology, SPPK No. 142/E5/PG.02.00PT/2022.

References

1. Bandyopadhyay D, Qureshi A, Ghosh S, Ashish K, Heise LR, Hajra A, Ghosh RK. Safety and Efficacy of Extremely Low LDL-Cholesterol Levels and Its Prospects in Hyperlipidemia Management. *J Lipids*. 2018; **2018**: 1-8.
2. Nelson RH. Hyperlipidemia as a Risk Factor for Cardiovascular Disease. *Prim Care*. 2013; **40**: 195-211.
3. Cruz PMR, Mo H, McConathy WJ, Sabnis N, Lacko AG. The Role of Cholesterol Metabolism and Cholesterol Transport in Carcinogenesis: A Review of Scientific Findings, Relevant to Future Cancer Therapeutics. *Front Pharmacol*. 2013; **4**: 119.
4. Golomb BA and Evans MA. Statin Adverse Effects: A Review of the Literature and Evidence for a Mitochondrial Mechanism. *Am J Cardiovasc Drugs*. 2008; **8**:373-418.
5. Ramkumar S, Raghunath A, Raghunath S. Statin Therapy: Review of Safety and Potential Side Effects. *Acta Cardiol Sin*. 2016; **32**: 631-639.
6. Sudirman S, Herpandi, Nopianti R, Lestari SD, Wasahla, Mareta H. Phenolic Contents, Tannin, Vitamin C, and Vitamin E of Water Lettuce (*Pistia stratiotes*). *Orient J Chem*. 2017; **33**: 3173-3176.
7. Herpandi, Lestari SD, Bastian, Sudirman S. Antioxidant Activity of the Fractions from Water Lettuce (*Pistia stratiotes*) Extract. *Food Res*. 2021; **5**: 451-455.
8. Islam B, Charu C, Adem A, Aburawi E, Ojha S. Insight into The Mechanism of Polyphenols on The Activity of HMGR by Molecular Docking. *Drug Des Dev Ther*. 2015; **9**: 4943-4951.
9. Zeka K, Ruparelia K, Arroo R, Budriesi R, Micucci M. Flavonoids and Their Metabolites: Prevention in Cardiovascular Diseases and Diabetes. *Dis*. 2017, **5**.
10. Dai J and Mumper RJ. Plant Phenolics: Extraction, Analysis and Their Antioxidant and Anticancer Properties. *Molecules*. 2010; **15**: 7313-7352.
11. Pérez-Magariño S, Ortega-Heras M, Cano-Mozo E. Optimization of a Solid-Phase Extraction Method Using Copolymer Sorbents for Isolation of Phenolic Compounds in Red Wines and Quantification by HPLC. *J Agric Food Chem*. 2008; **56**: 11560-11570.
12. Chandra S, Khan S, Avula B, Lata H, Yang MH, Elsohly MA, Khan IA. Assessment of Total Phenolic and Flavonoid Content, Antioxidant Properties, and Yield of Aero-ponically and Conventionally Grown Leafy Vegetables and Fruit Crops: A Comparative Study. *Evid-Based Compl Altern Med*. 2014; **2014**: 253875.
13. Barbosa-Pereira L, Pocheville A, Angulo I, Paseiro-Losada P, Cruz JM. Fractionation and Purification of Bioactive




Figure 1: Total polyphenol (A) and flavonoid (B) content of crude and purified extract of water lettuce (*Pistia stratiotes*) leaf. Data are shown as the mean \pm SD ($n=3$). Statistically significance at $*p<0.05$ versus purified extract.




Figure 2: Representative of HMG-CoA reductase activity and inhibitory activity of sample: (A) Blank (B) HMG-CoA reductase activity, (C) Crude extraction inhibitory activity, (D) Purified extract inhibitory activity, and (E) Pravastatin inhibitory activity (as a positive control). These activities were measured at 340 nm for 10 min by a spectrophotometer.




Figure 3: HMG-CoA reductase inhibitory activities of crude extract, purified extract, and pravastatin (as positive control). Data are shown as the mean \pm SD ($n=3$). Statistically significance at $*p<0.05$ versus purified extract.

Conclusion

The purified extract of water lettuce showed a high polyphenol and flavonoid contents when compared to the crude extract. The 3-hydroxy-3-methylglutaryl coenzyme-A reductase inhibitory activity of the purified extract was also higher when compared to crude extract. Therefore, the purified extract of polyphenol compounds from water

- Compounds Obtained from a Brewery Waste Stream. Biomed Res Int. 2013; **2013**: 1-11.
14. Yang Z, Tang H, Shao Q, Bilia A, Wang Y, Zhao X. Enrichment and Purification of the Bioactive Flavonoids from Flower of *Abelmoschus manihot* (L.) Medic using Macroporous Resins. Molecules. 2018; **23**: 2649.
 15. Lucci P, Saurina J, Núñez O. Trends in LC-MS and LC-HRMS Analysis and Characterization of Polyphenols in Food. Trends Anal Chem. 2017; **88**: 1-24.
 16. Souza JNS, Silva EM, da-Silva MN, Arruda MSP, Larondelle Y, Rogez H. Identification and Antioxidant Activity of Several Flavonoids of *Inga edulis* Leaves. J Braz Chem Soc. 2007; **18**: 1276-1280.
 17. Zhao H, Avena-Bustillos RJ, Wang SC. Extraction, Purification and *in vitro* Antioxidant Activity Evaluation of Phenolic Compounds in California Olive Pomace. Foods. 2022; **11**: 174.
 18. Baskaran G, Shukor MY, Salvamani S, Ahmad SA, Shaharuddin NA, Pattiram PD. HMG-CoA Reductase Inhibitory Activity and Phytocomponent Investigation of *Basella alba* Leaf Extract as a Treatment for Hypercholesterolemia. Drug Des Dev Ther. 2015; **9**: 509–517.
 19. Seenivasan A. Characterization, Modes of Synthesis, and Pleiotropic Effects of Hypocholesterolemic Compounds - A Review. Enzyme Inhib Med Chem. 2011; **4**: 23-32.
 20. Chen ZY, Jiao R, Ma KY. Cholesterol-Lowering Nutraceuticals and Functional Foods, J Agric Food Chem. 2008; **56**: 8761-8773.
 21. Friesen JA, Rodwell VW. The 3-hydroxy-3-methylglutaryl Coenzyme-A (HMG-CoA) Reductases. Genome Biol. 2004; **5**: 248..