

Tropical Journal of Natural Product Research

Available online at <https://www.tjnpr.org>

Original Research Article

Plectranthus scutellarioides Leaf Extract Protective Effects Against Isoniazid and Rifampicin-Induced Hepatotoxicity in Wistar Rats

Lusi P. Dwita¹, Ni Putu E. Hikmawanti^{2*}, Juhairoh Husniah¹, Muhammad Hazraj¹, Siti N. Bela¹, Eva Aryanti¹¹Department of Pharmacology, Faculty of Pharmacy and Sciences, Universitas Muhammadiyah Prof DR. HAMKA, East Jakarta, DKI Jakarta, 13460, Indonesia²Department of Pharmaceutical Biology, Faculty of Pharmacy and Sciences, Universitas Muhammadiyah Prof DR. HAMKA, East Jakarta, DKI Jakarta, 13460, Indonesia

ARTICLE INFO

Article history:

Received 12 August 2022

Revised 25 September 2022

Accepted 24 December 2022

Published online 01 January 2023

Copyright: © 2022 Dwita *et al.* This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Plectranthus scutellarioides has been studied to have antioxidant and anti-inflammatory activity which shows potential with hepatoprotection effect. This study determined the hepatoprotective activity of ethanol extract of *P. scutellarioides* leaves. The rats were divided into five groups: PSE (*Plectranthus scutellarioides* leaves extract) 150, 300, and 600 mg/kg, negative control, and positive control (Silymarin). All groups were given the test substances concurrently with isoniazid and rifampicin (50 mg/kg) p.o every day for 28 days. The results showed that the PSE could prevent liver damage as indicated by lower levels of total bilirubin, direct bilirubin, albumin, AST, and ALT than negative controls ($p < 0.005$). Histological observations of the liver confirmed these results, and this dose-dependent hepatoprotective activity was in line with the lipid peroxidation inhibition. *P. scutellarioides* extract at a dose of 600 mg/kg showed the best activity, comparable ($p > 0.05$) to Silymarin. The data suggest that the co-administration of *P. scutellarioides* leaf extract with isoniazid and rifampicin could prevent the liver injury caused by these drugs.

Keywords: Antioxidant, Flavonoids, Lamiaceae, Liver injury, Hepatoprotection, *Plectranthus scutellarioides*.

Introduction

Hepatoprotector protects liver cells from damage caused by hepatotoxic substances, such as drugs, chemical compounds, and toxic substances that can damage the liver.¹ A study on tuberculosis (TB) patients showed that combination therapy of isoniazid, rifampicin, and pyrazinamide is associated with liver injury severity.² The metabolism of isoniazid could produce hydrazine, which is toxic to the liver. Since rifampicin increases isoniazid metabolism, concomitant use of these drugs could worsen liver damage.³ Hepatoprotector concurrent use with TB drugs has been proven to be beneficial.

Several plants have been studied for their hepatoprotective activities, such as *Andrographis paniculata*, *Boerhaavia diffusa*, *Eclipta alba*, *Phyllanthus amarus*, *Picrorhiza kurroa*, *Mimosa pudica*, and *Silybum marianum*,⁴ where most of these plants also showed potential antioxidant activities. Antioxidant effects have been proven beneficial in preventing antitubercular hepatotoxicity risk,⁵ as well as the anti-inflammatory effect. *Plectranthus scutellarioides* leaves have shown potential for both activities.^{6,7} The flavonoid compounds of *P. scutellarioides* like apigenin 7-O-(3''-O-acetyl)- β -D-glucuronide, apigenin 5-O-(3''-O-acetyl)- β -D-glucuronide and rosmarinic acid, showed potent antioxidant activity.⁸ This study evaluated whether *P. scutellarioides* could reduce liver damage by isoniazid and rifampicin.

*Corresponding author. E mail: ermy0907@uhamka.ac.id
Tel: +62 85250874147

Citation: Dwita LP, Hikmawanti NPE, Husniah J, Hazraj M, Bela SN, Aryanti E. *Plectranthus scutellarioides* Leaf Extract Protective Effects Against Isoniazid and Rifampicin-Induced Hepatotoxicity in Wistar Rats. Trop J Nat Prod Res. 2022; 6(12):1936-1940. <http://www.doi.org/10.26538/tjnpr/v6i12.6>

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria

Materials and methods

P. scutellarioides leaves extraction

The leaves were obtained from Balitro, Indonesia and identified and authenticated as *Plectranthus scutellarioides* (L.) R. Br. from the family *Lamiaceae*. A total of 680 g of leaf powder was added with 7 L of 70% ethanol and macerated for 24 hours. The maceration was done seven times with new solvent. The filtrate was evaporated in a vacuum rotary evaporator (50°C) and then continued with a water bath (50°C) to obtain the thick extract.

Determination of *P. scutellarioides* leaves extract water content

Determination of extract characteristics and phytochemical content screening were done following the Indonesian Herb Pharmacopoeia. Ten grams of the thick extract was weighed and added with 200 mL of water-saturated toluene. The mixture was heated for approximately 2 hours, and then the receiver tube was cooled to room temperature. The volume of water was read and then calculated in % (v/b).

Determination of loss drying

The oven was set at 105°C, and the container was heated for 30 minutes, then weighed. The extract was weighed 1 g in the container, then put in an oven at 105°C until a fixed weight was obtained.

Phytochemical identification

The qualitative identification of phytochemical from *P. scutellarioides* Leaves Extract was conducted following the procedure in Indonesian Herbal Pharmacopoeia.⁹

Determination of total flavonoid levels

Serial concentrations of each extract were added with 3 ml of methanol, 0.2 ml of 10% AlCl₃, 0.2 ml of 1 M Na-acetate, and 10 ml of distilled water. The solution was incubated for 30 minutes, and then the absorbance was read at 415 nm using Spectrophotometer UV-Vis Seri UV-1601 (Shimadzu, Kyoto, Japan). The flavonoid concentration was determined with quercetin as the standard.

Total Flavonoid

$$= \frac{\text{Flavonoid concentration } (\mu\text{g/mL}) \times \text{sample volume (mL)} \times \text{dilution factor}}{\text{Extract weigh (g)}}$$

Animals and experimental design

Wistar rats (200-250 grams) were used for the test. The study protocol was approved by the Research Ethics Commission, Faculty of Medicine, University of Indonesia (404/UN2.F1/ETIK/PPM.00.02/2019). The rats were acclimatized for seven days before the experiment and kept under standard laboratory conditions (22–25 °C and a 12-h light/dark cycle), fed with a standard pellets diet and water ad libitum. The animals were divided into five groups: *Plectranthus scutellarioides* leaf extract (PSE) 150 mg/kg, 300 mg/kg, and 600 mg/kg, negative control and positive control (Silymarin) 100 mg/kg. All administration was done orally, 30 minutes prior to isoniazid and rifampicin. The TB drugs were given at the same dose, 50 mg/kg po. orally every day for 28 days. The next day, the rats were sacrificed by cervical dislocation.

Liver biochemical parameters

Blood samples were taken from the orbital sinuses on days 0, 14, and 28. The serum was used for total bilirubin, direct bilirubin, albumin, AST, and ALT levels. All parameters were determined using spectrophotometer clinical (MicroLab 300) with a reagent supplied by HUMAN Ltd., Germany.

Histopathological studies

The livers were fixed in 10% formalin for 24 hours. Then the histology preparations were prepared with Hematoxylin-Eosin staining. Histological observations, including the liver's pyknotic nucleus and Kupffer cells,¹⁰ were done in a 10-images view with 400× objective magnification.

Lipid peroxidation test

Lipid peroxidation of the liver was measured based on malondialdehyde (MDA) levels, with 1,1,3,3-tetra ethoxy propane (TEP) as standard. The liver was rinsed with phosphate buffer solution (PBS) pH 7.4 and dried. Then the liver was weighed and homogenized with PBS to make 20% homogenate. A total of 1 ml of homogenate was added with 0.5 ml of 20% trichloroacetic acid (TCA), then centrifuged at 3000 rpm for 10 minutes. 1 mL of the supernatant was added with 1 ml of 0.67% thiobarbiturate (TBA), heated at 100°C for 10 minutes, then cooled. The absorption was measured at 532 nm.

Statistical analysis

The data were analyzed with one-way ANOVA followed by Tukey's post-test.

Results and Discussion***Plectranthus scutellarioides* leaves extract characteristics**

Ethanol extract of *P. scutellarioides* leaves was obtained in the form of concentrated blackish-brown extract and a slightly bitter taste. The characteristics of *P. scutellarioides* leaves extract are presented in Table 1. Phytochemical screening showed that 70% ethanol extract of *P. scutellarioides* leaves contained flavonoids, phenols, saponins, and triterpenoids. The results showed that the total flavonoid level of 70% ethanol extract of *P. scutellarioides* leaves was 12.62 mg QE/g extract. Phytochemical screening showed that the *P. scutellarioides* leaves ethanol extract contained flavonoids, saponins, triterpenoids, and phenols. These compounds have been known to generate antioxidant activity, especially flavonoids.

Table 1: Characteristics of *P. scutellarioides* Leaves Extract

Characteristics	Results
Percentage of yield	18.41 %
Water content	6.71 %
Lost drying	7.84 %

The primary role of flavonoids as an antioxidant is to donate hydrogen atoms to free radicals that cause free radicals to become unreactive or neutral. The hepatoprotective activity of *P. scutellarioides* may be due to this flavonoid compound's presence. To confirm this hypothesis, we examined the total flavonoid content of the extract and the lipid peroxidation in the liver. The result showed that *P. scutellarioides* contains a high flavonoid (12.62 mg QE/g extract). Moreover, the lipid peroxidation test showed a significantly lower number of MDA compared to the negative control (Table 2).

Hepatoprotective Activity of *P. scutellarioides* leaves extract

Drug-induced liver injury (DILI) has been reported to occur in 19 per 10.000 people a year.¹² DILI can occur through an oxidative stress cascade, where the drug or its metabolites increase the reactive oxygen species (ROS), damaging the mitochondria and triggering immune reactions, thus initiating apoptosis or necrosis and causing cell death.¹³ This event eventually causes acute or chronic liver disease.¹⁴ The present study focuses on the approach to prevent liver injury against rifampicin and isoniazid toxicity. Both drugs have been known to cause hepatotoxicity in tuberculosis patients. It causes membrane damage, which increases liver enzymes and bilirubin in the systemic circulation, causing an imbalance in endogenous antioxidants and increased lipid peroxidation.¹⁵

This study used 50 mg/kg of isoniazid, and rifampicin to induce liver damage in rats. *P. scutellarioides* leave extract was given concurrently with the drugs for 28 days. The hepatoprotective activities were examined by measuring the levels of serum total bilirubin, direct bilirubin, AST, ALT, albumin, liver lipid peroxidation, and histology examination (Figure 1 A – E).

The result showed that the administration of isoniazid and rifampicin for 28 days increased total bilirubin levels by 56.89%, direct bilirubin by 61.37%, and lowered albumin levels by 40.86%. on the other hand, concurrent administration of PSE with the drugs significantly lowered total bilirubin, direct bilirubin, and higher albumin levels than the negative control (p <0.05). The best result seen in PSE 600mg/kg group, where the total bilirubin and direct bilirubin levels, respectively, were 53.45% and 64.98% lower than negative control on day 28, while albumin levels remained stable during the treatment.

Isoniazid and rifampicin caused liver injury, as indicated by the drastic change in all hepatic biochemical parameters since day 14. On the other hand, PSE groups showed improvement in all parameters, except for 150 mg/kg. The results show that PSE treatment groups showed a significant decreased in total bilirubin levels and direct bilirubin, while the albumin level was higher (p <0.05) than negative controls at 300 and 600 mg/kg doses. The same trend was seen in the ALT and AST levels. It was also observed that 600 mg/kg of PSE resulted in stable hepatic parameters and almost no chance compared to day 0 (Figure 2A - C).

The liver enzyme levels in the systemic circulation are indicative of liver cell damage, whereas bilirubin and albumin levels are associated with liver function. Measurement of bilirubin levels in this study was done by the Jendrassik-Grof method, where the bilirubin reacts with Diazotized Sulfanilic Acid (DSA) and forms azobilirubin. Total bilirubin consists of direct (conjugated) and indirect (unconjugated) bilirubin, where glucuronic acid, with the help of uridine diphosphate, conjugates with the bilirubin to form direct bilirubin.¹⁶

Oxidative stress due to the use of isoniazid and rifampicin also results in leakage of the liver cell membrane and causes the release of transaminase enzymes into the circulation, such as ALT and AST.¹⁷ ALT is present in the cytosol of the liver, while AST is found in the mitochondria of the liver, heart, kidneys, and brain. The damage to these tissues causes an increase in serum ALT and AST levels.¹⁸ PSE group at doses 300 and 600 mg/kg showed a significantly lower level of both AST and ALT than the negative control, indicating its activity in preventing liver cell damage. However, the lowest dose of PSE showed an insignificant effect.

Based on the data, *P. scutellarioides* leaves extract showed hepatoprotective activity in dose depending manner, where the highest dose showed the best activity comparable to Legalon® (equivalent to silymarin 272.15 mg/kg rats BW). Silymarin is a natural compound derived from *Silybum marianum*.¹⁹ It could prevent damage caused by

tuberculosis drugs through a mechanism of reducing oxidative stress and lipid peroxidation.²⁰ Silymarin stabilizes the reactive oxygen species (ROS), especially in the liver, and therefore is usually used in hepatitis, hepatic cirrhosis, and other liver disorders.²¹ Histological observations of liver cells show that isoniazid and rifampicin cause inflammation and cell death (necrosis), which is in line with other studies,¹¹ where these TB drugs induce inflammation characterized by an increase in the number of Kupffer cells. The drugs also cause cell necrosis, as indicated by a shrinking and blackish cell nucleus (pyknotic nucleus).¹¹ PSE co-administration with isoniazid and rifampicin showed a noticeable histology difference compared to the negative control, especially at 600 mg/kg. PSE could reduce damage in the liver shown by Kupffer cells and the pyknotic nuclei in

liver cells (Table 2). Figure 2 depicts a similar observation in Legalon® and PSE 600 mg/kg with normal control, suggesting the hepatoprotective effect of these groups.

In the present study, histological observations were carried out by counting the number of pyknotic nuclei and kupffer cells in liver cells (Table 2). Isoniazid and rifampicin cause the accumulation of toxic substances in liver cells that cause cell death. Microscopically visible changes in the cell nucleus seen as wrinkled or shrunk, denser and dark, with irregular cell boundaries (pyknotic nucleus). Whereas the number of Kupffer cells, which are macrophages in the liver, are increased, showing an inflammatory response.¹¹ Kupffer cells have an oval shape on the surface of cells in the sinusoid.²² The identical observation was also seen in humans.




Figure 1: Hepatoprotective effect of *P. scutellarioides* leaves extract day 0 (before treatment), 14 and 28, (A) aspartate aminotransferase (AST), (B) alanine aminotransferase (ALT), (C) Albumin Levels (D) Direct bilirubin, (E) Total bilirubin. The means in each day that do not share a letter are significantly different ($p < 0.05$)




Figure 2: Histological Observation of Livers (A) Normal control (B) Negative control (C) Legalon® (D) PSE 150 mg/kg (E) PSE 300 mg/kg (F) PSE 600 mg/kg (1) Pyknotic nucleus (2) Kupffer Cells (400x), PSE (*Plectranthus scutellarioides* Leaves extract).

Table 2: The Histological and Chemical Examination of The Rats Liver

Groups	Pyknotic nuclei X ± SD, n=4	Kupffer cells X ± SD, n=4	MDA X ± SD, n=4
PSE 150mg/kg	113.0 ± 11.2 ^d	75.32 ± 6.21 ^d	19.51 ± 0.42
PSE 300 mg/kg	93.00 ± 4.69 ^e	58.45 ± 7.84 ^e	17.70 ± 0.45 ^b
PSE 600 mg/kg	71.55 ± 11.1 ^c	41.25 ± 3.10 ^c	15.63 ± 0.69 ^d
Legalon®	62.80 ± 8.03 ^c	31.70 ± 2.70	13.04 ± 0.83
Negative control	136.2 ± 7.99 ^b	88.00 ± 7.25	21.90 ± 0.84 ^e
Normal control	47.90 ± 7.95 ^a	21.80 ± 3.29 ^a	12.74 ± 1.07 ^c

The same subscript in each column shows comparable results ($p > 0.05$), PSE (*Plectranthus scutellarioides* Leaves extract)

Liver tissue biopsy in patients taking these drugs shows an increase in inflammatory cells, especially histiocytes and neutrophils.²³ The number of pyknotic nuclei and kupffer cells in the group that received PSE showed mild liver damage compared to the negative control group, especially at a dose of 600 mg/kg.

Studies have shown several potential plants as hepatoprotector. Another species of *Plectranthus*, *P. amboinicus* also showed hepatoprotective effect at 600 mg/kg doses in paracetamol-induced rats.²⁴ *Plectranthus* genus exhibits hepatoprotective activity at the same dose, despite the different drugs used to induce liver injury. Despite PSE's success in preventing liver damage, satisfactory results were only seen at high doses. Compared to other studies, such as *Telfairia occidentalis* at a dose of 500mg/kg²⁵ and *Alpinia Officinarum* at a dose of 400 mg/kg,²⁶ showed a lower dose as hepatoprotector against rifampicin and isoniazid. The biological activity of *P. scutellarioides* leaves extract was due to the complexity of the chemical compounds contained therein. The fractions/isolates of this plant should be further studied so that the development of *P. scutellarioides* leaves as hepatoprotector can be optimal.

Conclusion

Based on liver enzyme levels and liver function tests, PSE showed beneficial effects in preventing liver damage due to isoniazid and rifampicin hepatotoxicity. It can be concluded that PSE has hepatoprotective potential, especially at a dose of 600 mg/kg.

Conflict of Interest

The authors declare no conflict of interest.

Authors' Declaration

The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.

Acknowledgments

This work was supported by Lembaga Penelitian dan Pengembangan Universitas Muhammadiyah Prof DR HAMKA.

References

- Dong Y, Huang J, Lin X, Zhang S, Jiao Y, Liang T, Chen Z, Huang R. Hepatoprotective effects of Yulangsan polysaccharide against isoniazid and rifampicin-induced liver injury in mice. *J Ethnopharmacol.* 2014 Feb 27;152(1):201–6.
- Zhao H, Wang Y, Zhang T, Wang Q, Xie W. Drug-Induced Liver Injury from Anti-Tuberculosis Treatment: A Retrospective Cohort Study. *Med Sci Monit.* 2020;26:e920350.

3. Ramappa V, Aithal GP. Hepatotoxicity Related to Anti-tuberculosis Drugs: Mechanisms and Management. *J Clin Exp Hepatol*. 2013;3(1):37–49.
4. Shakya AK. Drug-induced hepatotoxicity and hepatoprotective medicinal plants: A review. *Indian J Pharm Educ Res*. 2020;54(2):234–50.
5. Basheer AS, Siddiqui A, Paudel YN, Hassan MQ, Imran M, Najmi AK, Akhtar M. Hepatoprotective and antioxidant effects of fish oil on isoniazid-rifampin induced hepatotoxicity in rats. *PharmaNutrition* [Internet]. 2017;5(1):29–33. Available from: <http://dx.doi.org/10.1016/j.phanu.2017.01.002>
6. Moelyono MW, Uswatun A, Rochjana H, Diantini A, Mus I, Sumiwi SA, Iskandar Y, Susilawati, Y. Antioxidant Activity of *Plectranthus scutellarioides* (L.) Leaves. *J Farm Indones*. 2016;8(1):271–6.
7. Levita J, Sumiwi SA, Pratiwi TI, Ilham E, Sidiq SP, Moektiwardoyo M. Pharmacological Activities of *Plectranthus scutellarioides* (L.) R.Br. Leaves Extract on Cyclooxygenase and Xanthine Oxidase Enzymes. *J Med Plants Res*. 2016;10(20):261–9.
8. Kubínová R, Gazdová M, Hanáková Z, Jurkaninová S, Acqua SD, Cvačka J, Humpa O. South African Journal of Botany New diterpenoid glucoside and flavonoids from *Plectranthus scutellarioides* (L.) R. Br. 2019;120:286–90.
9. Indonesia M of HR of. *Farmakope Herbal Indonesia* (Indonesian Herb Pharmacopoeia). Jakarta: Ministry of Health Republic of Indonesia; 2008. 221 p.
10. Kiernan JA. *Histological and Histochemical Methods*. 5th ed. Chatham, UK: Scion; 2015.
11. Siddique W, Rashid N, Warraich NY, Asghar A, Talat H, Zia AE, Fatima A. Evaluation of histopathological changes in isoniazid and rifampicin induced liver injury and protective role of zinc sulfate in Albino rats. *Pakistan J Med Heal Sci*. 2019;13(4):809–16.
12. Björnsson ES. Epidemiology, Predisposing Factors, and Outcomes of Drug-Induced Liver Injury. Vol. 24, *Clinics in Liver Disease*. W.B. Saunders; 2020. p. 1–10.
13. Teschke R, Danan G. Drug Induced Liver Injury: Mechanisms, Diagnosis, and Clinical Management. In: *Liver Diseases*. Springer International Publishing; 2020. p. 95–105.
14. Kaplowitz N. Drug-Induced Liver Injury. *Clin Infect Dis*. 2004;38(Suppl 2):44–8.
15. Tasduq SA, Kaiser P, Sharma SC, Johri RK. Potentiation of isoniazid-induced liver toxicity by rifampicin in a combinational therapy of antitubercular drugs (rifampicin, isoniazid and pyrazinamide) in Wistar rats: A toxicity profile study. *Hepatol Res* [Internet]. 2007 Oct 1 [cited 2020 Apr 15];37(10):845–53. Available from: <http://doi.wiley.com/10.1111/j.1872-034X.2007.00129.x>
16. Kumar V, Abbas AK, Aster JC. *Robbins, Basic Pathology*. The British Journal of Psychiatry. Elsevier Inc; 2018.
17. Lee EH, Baek SY, Park JY, Kim YW. Rifampicin activates AMPK and alleviates oxidative stress in the liver as mediated with Nrf2 signaling. *Chem Biol Interact* [Internet]. 2020;315(108889):1–8. Available from: <https://doi.org/10.1016/j.cbi.2019.108889>
18. Mccorry LK, Zdanowicz MM, Gonnella CY. *Essentials of Human Physiology and Pathophysiology for Pharmacy and Allied Health*. The British Journal of Psychiatry. Routledge, Taylor & Francis Group; 2019.
19. Vargas-Mendoza N, Madrigal-Santillán E, Morales-González Á, Esquivel-Soto J, Esquivel-Chirino C, García-Luna y González-Rubio MG, et al. Hepatoprotective effect of silymarin. *World J Hepatol*. 2014;6(3):144–9.
20. Tao L, Qu X, Zhang Y, Song Y, Zhang SX. Prophylactic Therapy of Silymarin (Milk Thistle) on Antituberculosis Drug-Induced Liver Injury: A Meta-Analysis of Randomized Controlled Trials. *Can J Gastroenterol Hepatol*. 2019;1–11.
21. Karimi G, Vahabzadeh M, Lari P, Rashedinia M, Moshiri M. “Silymarin”, a Promising Pharmacological Agent for Treatment of Diseases. 2011;14(4):308–17.
22. Junqueira L, Carneiro J. *Basic Histology: Text & Atlas*. The McGraw-Hill Companies. 2007.
23. Ibrahim A, Payus AO. “A Double-Edged Sword”: A Case Report on Liver Injury Side-Effect of Anti-Tuberculosis Drug on Liver Tuberculosis Patient”. *Borneo J Med Sci* [Internet]. 2018;14(1):49–53.
24. Shenoy S, Kumar H, Thashma, Nayak V, Prabhu K, Pai P, Warriar I, Somayaji, Madhav V, Bairy KL, Kishore A. Hepatoprotective activity of *Plectranthus amnoenicus* against paracetamol induced hepatotoxicity in rats. Vol. 1, *International Journal of Pharmacology and Clinical Sciences*. 2012. p. 32–8. Available from: <https://www.ijpcs.net/article/46>
25. Nwidu LL, Oboma YI. *Telfairia occidentalis* (Cucurbitaceae) pulp extract mitigates rifampicin-isoniazid-induced hepatotoxicity in an in vivo rat model of oxidative stress. *J Integr Med* [Internet]. 2019;17(1):46–56. Available from: <https://doi.org/10.1016/j.joim.2018.11.008>
26. Eram S, Mujahid M, Bagga P, Ahsan F, Rahman MA. Hepatoprotective Evaluation of Galanga (*Alpinia Officinarum*) Rhizome Extract against Antitubercular Drugs Induced Hepatotoxicity in Rats. *J Herbs, Spices Med Plants*. 2020;26(2):113–25.